

Cluster formation in complex multi-scale systems

J. D. Gibbon

Department of Mathematics, Imperial College London, London SW7 2AZ, UK

E. S. Titi

Department of Computer Science and Applied Mathematics,

Weizmann Institute of Science,

P.O. Box 26, Rehovot, 76100 Israel

and

Department of Mathematics and Department of Mechanical and Aerospace Engineering,

University of California, Irvine, CA 92697-3875, USA

(Dated: February 2, 2008)

Based on the competition between members of a hierarchy of length scales in complex multi-scale systems, it is shown how clustering of active quantities into concentrated sets, like bubbles in a Swiss cheese, is a generic property that dominates the intermittent structure. The halo-like surfaces of these clusters have scaling exponents lower than that of their kernels, which can be as high as the domain dimension. Examples include spots in fluid turbulence and droplets in spin-glasses.

PACS numbers: 47.10.+g, 47.27.Ak, 89.75.-k, 05.45.a

Keywords: Clustering, intermittency, fluid turbulence, complex systems, multi-scale, spin glass.

It has long been recognized that active quantities in complex systems of many types are not distributed evenly across a domain but cluster strongly into irregular bubbles, as in a Swiss cheese. The nomenclature, the nature and shape of the bubbles, and the physics in each subject is substantially different: spottiness in high Reynolds number fluid turbulence [1, 2, 3, 4, 5, 6, 7, 8, 9] and boundary layers [10]; droplet formation in spin-glasses [11, 12, 13]; clustering behaviour in networks [14, 15]; the preferential concentration of inertial particles [16, 17, 18, 19, 20] with applications to rain initiation by cloud turbulence [21]; clustering of luminous matter [22, 23, 24, 25] and magnetic bubbles in astro-physics [26], are some examples. These clusters display strong features whose typical length scales are much shorter than their averages, thus raising the question of the nature of the interface between them and the surrounding longer scale regions. For instance, in spin glasses the ‘surface’ of the droplets has a fractal-like structure whereas the droplets themselves have the full domain dimension [13]. In fluid turbulence the concentrated sets on which vorticity accumulates are tubes and sheets, although the fractal nature of these is unclear. These sets dominate the associated Fourier spectra which display a spikiness that is the hallmark of what is usually referred to as intermittency [1, 2, 3, 4, 5, 6, 7, 8, 9]. The ubiquity of this irregular bubble-like topology suggests the existence of a set of underlying organizing principles in complex multi-scale systems. *Using simple but broadly applicable mathematical ideas, this paper will demonstrate that the dominant physical principle behind clustering is the existence of a hierarchy of length scales whose members are in competition.*

Consider a d -dimensional system whose smallest characteristic (integral) scale L is such that the system is statistically homogeneous on boxes $\Omega = [0, L]^d$. Moreover,




FIG. 1: An illustrative slice through Ω for one value of n : the black kernels are surrounded by green(gray) halos. Very small-scale behaviour concentrates on the black & green(gray) regions which constitute the set $\mathcal{A}_n^+$ ($\mathcal{L}_n \kappa_n > 1$). The halos have scaling exponents lower than those of the black kernels.

it is endowed with the following two properties. Firstly, at each point $x \in \Omega$, it possesses an ordered set of length scales $\ell_n = \ell_n(x)$ associated with a hierarchy of features labelled by $n \geq 2$

$$L > \ell_1 \geq \ell_2 \geq \dots \geq \ell_n \geq \ell_{n+1} \dots \quad (1)$$

The ℓ_n could be thought of as an ordered set of correlation or coherence lengths; their inverses $\kappa_n(x) = \ell_n^{-1}(x)$ clearly obey $1 < L\kappa_n \leq L\kappa_{n+1}$. The second assumption is that the ensemble averages of the $L\kappa_n(x)$ are bounded above by some ordered, positive parameters of the system satisfying $1 < R_n \leq R_{n+1}$

$$1 < L \langle \kappa_n \rangle \leq R_n. \quad (2)$$

The ensemble average $\langle \cdot \rangle$ is a spatial average with respect to the Lebesgue measure over Ω . Thus, while the ordering of the $\ell_n(x)$ must be respected at each point, the ℓ_n themselves could be quite rough; e.g. they could

consist of a series of step functions. If they become very small near points x^* then they must obey $\ell_n > O(r^{d-\varepsilon})$ ($r = |x - x^*|$ and $\varepsilon > 0$) so as not to violate (2).

Following an idea used in ref. [27], consider the real arbitrary parameters $0 < \mu < 1$ and $0 < \alpha < 1$ such that $\mu + \alpha = 1$. Use Hölder's inequality, $|AB| \leq \frac{1}{p}|A|^p + \frac{1}{q}|B|^q$ where $\frac{1}{p} + \frac{1}{q} = 1$, with $p = \mu^{-1}$ and $q = \alpha^{-1}$

$$\begin{aligned} \langle \kappa_n^\alpha \rangle &\leq \langle \kappa_{n+1}^\alpha \rangle = \left\langle \left(\frac{\kappa_{n+1}}{\kappa_n} \right)^\alpha \kappa_n^\alpha \right\rangle \\ &\leq \left\langle \left(\frac{\kappa_{n+1}}{\kappa_n} \right)^{\alpha/\mu} \right\rangle^\mu \langle \kappa_n \rangle^\alpha. \end{aligned} \quad (3)$$

Re-arranging and factoring out a term $\langle \kappa_n^\alpha \rangle$ gives

$$\left\langle \left(\frac{\kappa_{n+1}}{\kappa_n} \right)^{\alpha/\mu} \right\rangle \geq \langle \kappa_n^\alpha \rangle \left(\frac{\langle \kappa_n^\alpha \rangle}{\langle \kappa_n \rangle} \right)^{\alpha/\mu}. \quad (4)$$

Lower bounds on the ratio $\langle \kappa_n^\alpha \rangle / \langle \kappa_n \rangle$ can be found from (2) thereby turning (4) into

$$\left\langle \left(\frac{\kappa_{n+1}}{\kappa_n} \right)^{\alpha/\mu} \right\rangle - [(L\kappa_n)^\mu R_n^{-1}]^{\alpha/\mu} \geq 0. \quad (5)$$

While it is possible that the integrand in (5) could be positive everywhere in Ω , this cannot be assumed; the generic case is that the integrand could take either sign[40]. With the definition $\mathcal{L}_n = L R_n^{-1/\mu}$ we have the pair of inequalities

$$\frac{\kappa_{n+1}}{\kappa_n} \geq (\mathcal{L}_n \kappa_n)^\mu, \quad (6)$$

for which $\geq$ is valid on regions where the integrand is positive, designated as *good regions*, and negative ($<$) on *bad regions*. The term $(\mathcal{L}_n \kappa_n)^\mu$ on the right hand side of (6) remarkably contains the arbitrary parameter μ which lies in the range $0 < \mu < 1$. Its existence is important because the ordering in (1) makes it clear from (6) that everywhere within the bad regions ($<$) there are large lower bounds on κ_n with exponents containing $1/\mu$

$$\mathcal{L}_n \kappa_n > 1, \quad \Rightarrow \quad L\kappa_n > R_n^{1/\mu}. \quad (7)$$

Let $\mathcal{A}_n^+$ be the set on which $\mathcal{L}_n \kappa_n > 1$ and $\mathcal{A}_n^-$ the set on which $\mathcal{L}_n \kappa_n \leq 1$. Then all the bad regions ($<$), designated by the clusters of black kernels in Figure 1, lie in $\mathcal{A}_n^+$. The green/gray halos also lie in $\mathcal{A}_n^+$, and correspond to those parts of the good regions ($\geq$) neighbouring the bad. It is in these halos where the lower bound $(\mathcal{L}_n \kappa_n)^\mu$ becomes operative. The white areas of Figure 1 correspond to $\mathcal{A}_n^-$ in which the κ_n can be randomly distributed subject to their ordering in (1). It is clear from (5) that the existence and location of the clusters may differ for each n . A physical picture that displays all clusters for every n would be the union $\mathcal{A}^+ = \cup \mathcal{A}_n^+$.

To show that the volume $\mathcal{V}_n^+$ of $\mathcal{A}_n^+$ comprises a small part of Ω , Chebychev's inequality relates the normalized

Lebesgue measure $m(\mathcal{A}_n^+)$ to the integral of $L\kappa_n$ over $\mathcal{A}_n^+$

$$\int_{\mathcal{A}_n^+} L\kappa_n dm \geq m(\mathcal{A}_n^+) R_n^{1/\mu} = L^{-d} \mathcal{V}_n^+ R_n^{1/\mu}. \quad (8)$$

Together with the relation $\int_{\mathcal{A}_n^+} L\kappa_n dm \leq \langle L\kappa_n \rangle \leq R_n$ we have

$$m(\mathcal{A}_n^+) \leq R_n^{-\frac{1}{\mu}+1}. \quad (9)$$

Hence $m(\mathcal{A}_n^+)$ is significantly smaller than unity and decreases as R_n increases. Thus $\mathcal{A}_n^+$ can fill, at most, a small fraction of Ω . With such sparse information it is difficult to estimate the Hausdorff or the fractal dimensions of $\mathcal{A}_n^+$, but it is still possible to estimate scaling exponents [28]. This entails making a third assumption of self-similarity to estimate the smallest number of balls $\mathcal{N}_n^+$ of radius λ_n^+ needed to cover $\mathcal{A}_n^+$. Defining λ_n^+ as

$$(\lambda_n^+)^{-1} \equiv k_n^+ = \langle \kappa_n^p \rangle^{1/p}, \quad (10)$$

for some $p > 1$, it is clear that k_n^+ cannot be large enough when $p = 1$ because of (2). However, any value[41] of $p \gg 1$ will do that makes k_n^+ large enough to be a member of $\mathcal{A}_n^+$. The simplest and worst estimate would be to write

$$\mathcal{N}_n^+ \sim (L/\lambda_n^+)^d. \quad (11)$$

Inequality (9), however, shows that $\mathcal{A}_n^+$ occupies only a small fraction of Ω . A multiplicative factor of $m(\mathcal{A}_n^+)$ is introduced thus

$$\mathcal{N}_n^+ \sim m(\mathcal{A}_n^+) \left(\frac{L}{\lambda_n^+} \right)^d = m(\mathcal{A}_n^+) (Lk_n^+)^d. \quad (12)$$

Instead of using (9) to estimate $m(\mathcal{A}_n^+)$, an assumption of self-similar scaling is introduced that requires that the change in volume of the balls with respect to n should scale as $\mathcal{V}_n^+$ (the volume of $\mathcal{A}_n^+$) scales to L^d . Thus

$$m(\mathcal{A}_n^+) \sim \frac{\mathcal{V}_n^+}{L^d} \sim \left(\frac{\lambda_{n+1}^+}{\lambda_n^+} \right)^d. \quad (13)$$

We observe that the definition of the set $\mathcal{A}_n^+$ in principle involves the length scales L and λ_n^+ , but not overtly λ_{n+1}^+ . Yet the good and bad sets involve all three scales; L , λ_n^+ and λ_{n+1}^+ . The self-similarity assumption (13) is an assumption about the nature of the set $\mathcal{A}_n^+$ that relates successive length scales λ_n^+ and λ_{n+1}^+ in an *ad hoc*, yet reasonable, fashion. Using (13) in (12) we have

$$\mathcal{N}_n^+ \sim \left(\frac{\lambda_{n+1}^+}{\lambda_n^+} \right)^d \left(\frac{L}{\lambda_n^+} \right)^d = \frac{(\mathcal{L}_n k_n^+)^{2d}}{(\mathcal{L}_n k_{n+1}^+)^d} R_n^{d/\mu}. \quad (14)$$

From these, two estimates for $\mathcal{N}_n^+$ emerge, one each for the green/gray halo and black kernel regions of Figure 1, whose scaling exponents[42] are independent of p

$$\mathcal{N}_n^+ \lesssim \begin{cases} (\mathcal{L}_n k_n^+)^{d(1-\mu)} R_n^{d/\mu} & \text{(green/gray halo)} \\ (\mathcal{L}_n k_n^+)^d R_n^{d/\mu} & \text{(black kernel)} \end{cases} \quad (15)$$

TABLE I: Summary of conclusions regarding the sets $\mathcal{A}_n^\pm$ and the coloured regions in Figure 1.

Figure 1	black	green(gray)	white
Set	$\mathcal{A}_n^+$	$\mathcal{A}_n^+$	$\mathcal{A}_n^-$
Inequality (6)	$< \text{(bad)}$	$\geq \text{(good)}$	$\geq \text{(good)}$
Exponent	$\leq d$	$\leq d(1 - \mu)$	$= d$

For the former, the $>$ direction of the inequality in (6) has been used together with a simple Hölder inequality

$$\left\langle \kappa_n^{p(1+\mu)} \right\rangle^{1/p} \geq \langle \kappa_n^p \rangle^{(1+\mu)/p} = (k_n^+)^{1+\mu}. \quad (16)$$

whereas for the latter $\kappa_n \leq \kappa_{n+1}$ has been used. In contrast, without any evidence of contraction of volume, the formula corresponding to (12) for $\mathcal{N}_n^-$ is

$$\mathcal{N}_n^- \sim \left(\frac{L}{\lambda_n^-} \right)^d = (\mathcal{L}_n k_n^-)^d R_n^{d/\mu}, \quad (17)$$

where k_n^- satisfies $\mathcal{L}_n \kappa_n^- \leq 1$. The uniform scaling exponents in (15) are bounded by

$$\mathcal{D}_{n,\text{halo}}^+ \leq d(1 - \mu) \quad \mathcal{D}_{n,\text{ker}}^+ \leq d \quad (18)$$

whereas $\mathcal{D}^- = d$ from (17). The coefficients $R_n^{d/\mu}$ in (14) to (17) reflect the fact that this effect is taking place only at length scales smaller than $LR_n^{-1/\mu}$. The green/gray halo clearly plays the role of an interface of small but finite thickness between the d -dimensional (white) outer region and the (black) inner kernel whose dimension can be as high as d but could be less. When $\mathcal{D}_{n,\text{ker}}^+$ saturates its upper bound we have

$$\mathcal{D}_{n,\text{halo}}^+ \leq d(1 - \mu) < \mathcal{D}_{n,\text{ker}}^+ = d. \quad (19)$$

For the green/gray region to have an exponent at least $d - 1$ (a surface), μ would lie in the range $0 < \mu \leq 1/d$. Without equations of motion, a numerical experiment would be necessary to estimate the R_n by finding the maximum value of the ensemble average $\langle \kappa_n \rangle$. In principle μ could then be found from numerical estimates of $\ell_n^{\text{crit}} \sim LR_n^{-1/\mu}$ within the black kernels although if the κ_n take very large values there it might not be possible to achieve resolution. μ itself may have upper and lower bounds that are themselves n -dependent, as in ref. [27].

We now proceed to discuss some examples. The first ideas on clustering came more than half a century ago from Batchelor and Townsend [2] who observed intermittent behaviour in their high Reynolds number flow experiments, closely followed by observations in boundary layers by Emmons [10]. Batchelor and Townsend called this phenomenon ‘spottiness’ and suggested that the energy associated with the small scale components is distributed unevenly in space and roughly confined to regions which concomitantly become smaller with eddy size [29]. Mandelbrot then suggested that these clustered

sets on which energy dissipation is the greatest might be fractal in nature [30]. In measurements of the energy dissipation rate in the atmospheric surface layer, Meneveau and Sreenivasan interpreted the intermittent nature of their signals in terms of multi-fractals [3]; a newer generation of experiments measuring intense dissipation in turbulent flows have been pursued by Zeff *et al* [4]. Sreenivasan and Bershadskii have recently suggested that the clustering of high frequencies in a turbulent signal can be characterized by a scaling exponent [5].

The extremely rapid time evolution of sets of high vorticity or strain in fluid turbulence is an important issue; many computations exist showing how these take on the nature of quasi-one-dimensional tubes and quasi-two-dimensional sheets which have short lifetimes [6, 9]. An alternative to studying the problem in a statistical manner is to include time in the ensemble average $\langle \cdot \rangle$, in which case the semi-infinite nature of the time-axis suggests a different measure might be necessary [43]. With specific reference to the Navier-Stokes equations, analysis is not advanced enough to deal with the full space-time equations (except see ref. [31]); conventional methods of analysis use Sobolev norms to L^2 -average over space and remove the pressure [6, 32, 33] leaving only time. In ref. [27] a hierarchy of κ_n have been constructed which are comprised of ratios of norms (of derivatives of order n) and therefore functions of time only; thus the clusters of Figure 1 are merely gaps in the time-axis. It is then necessary to prove that they are finite in width and decreasing with increasing Reynolds number. This involves finding bounds on μ .

The second example is that of the low-temperature phase of spin glasses [34, 35]. Our conclusions regarding the fundamental role played by the competition between members of a hierarchy of length scales is consistent with the observation of ultrametricity in spin-glasses, a term that is used to denote the presence of a hierarchy of scales [35, 36]. This has been observed in computations on the low-temperature spin glass phases of the Sherrington-Kirkpatrick [34, 36, 37] and Edwards-Anderson models [38], as well as in dynamic phenomena in complexity [39]. The results of this paper, and particularly with reference to (19), are consistent with the droplet theory [11, 12, 13] where the kernel of the droplet is of full dimension d but its surrounding ‘surface’ has a scaling exponent $< d$. In fact, Palassini and Young [13] have shown numerically that $\mathcal{D}_{\text{halo}}^+ = 2.58 \pm 0.02$ when $d = 3$ and $\mathcal{D}_{\text{halo}}^+ = 2.77 \pm 0.02$ when $d = 4$.

In conclusion, we have shown that in a system endowed with a competitive hierarchy of correlation lengths, a clustering effect ensues in which length scales smaller than a critical value, and much smaller than the ensemble average scale, aggregate into small intense regions. The kernels of these intense regions are surrounded by halos that have scaling exponents smaller than of the domain dimension d . We have expectations that this idea of competition between scales may be a useful paradigm in explaining the behaviour of multi-scale systems.

Acknowledgments

We wish to acknowledge discussions with Steve Cowley, Charles Doering, Darryl Holm, Roy Jacobs, Robert Kerr, Michael Moore, Maya Paczusi, Andrew Parry, Greg Pavliotis, Jaroslav Stark and Christos Vassilicos. J.D.G. would like to thank the Isaiah Berlin Foundation for travel support and the hospitality of the Faculty of

Mathematics and Computer Science of the Weizmann Institute of Science where this work was begun. The work of E.S.T. was supported in part by the NSF grant number DMS-0204794, an MAOF Fellowship of the Israeli Council of Higher Education, the USA Department of Energy under contract number W-7405-ENG-36 and the ASCR Program in Applied Mathematical Sciences.

-
- [1] U. Frisch *Turbulence: The legacy of A. N. Kolmogorov* (Cambridge University Press, Cambridge 1995).
 - [2] G. K. Batchelor and A. Townsend, Proc. R. Soc. Lond. A, **199**, 238 (1949).
 - [3] C. Meneveau and K. Sreenivasan, J. Fluid Mech. **224**, 429 (1991).
 - [4] B. W. Zeff, D. D. Lanterman, R. McAllister, R. Roy, E. J. Kostelich and D. P. Lathrop, Nature, **421**, 146 (2003).
 - [5] K. Sreenivasan and A. Bershadskii, *Scaling clustering in turbulent signals*, preprint (2005).
 - [6] A. J. Majda and A. Bertozzi, *Vorticity and Incompressible Flow* (Cambridge University Press, Cambridge 2002).
 - [7] A. J. Majda, *Introduction to P.D.E.s and Waves for the Atmosphere and Ocean*, Courant Lecture Notes **9**, (AMS & Courant Institute of Mathematical Sciences 2002).
 - [8] R. M. Kerr, J. Fluid Mech., **153**, 31 (1985).
 - [9] A. Vincent and M. Meneguzzi, J. Fluid Mech., **225**, 245 (1994).
 - [10] H. W. Emmons, J. Aero Sci. **18**, 490 (1951).
 - [11] D. Fisher and D. A. Huse, Phys. Rev. Lett. **56**, 1601 (1986).
 - [12] A. J. Bray and M. A. Moore, Phys. Rev. Lett. **58**, 57 (1987).
 - [13] M. Palassini and A. P. Young, Phys. Rev. Lett. **85**, 3017 (2000).
 - [14] R. Albert and A.-L. Barabási, Rev. Mod. Phys. **74**, 47 (2002).
 - [15] G. B. West, J. H. Brown and B. J. Enquist, Science, **284**, 1677, (1999).
 - [16] J. K. Eaton and J. R. Fessler, Int. J. Multiphase Flow **20**, 169, (1994).
 - [17] H. Sigurgeirsson and A. M. Stuart, Phys. Fluids, **14**, 4352, (2002).
 - [18] J. Bec, *Multifractal concentrations of inertial particles in smooth random flows*, (2004). preprint
 - [19] J. Bec and A. Celani and M. Cencini and S. Musacchio, *Clustering and collisions of heavy particles in random smooth flows*, preprint (2004).
 - [20] D. D. Holm and V. Putkaradze, *Aggregation of finite particles with variable mobility*, <http://arxiv.org/abs/nlin.PS/0501009>.
 - [21] G. Falkovich, A. Fouxon and M. G. Stepanov, Nature, **419**, 151, (2002).
 - [22] P. Bak and K. Chen, Phys. Rev. Lett. **86**, 4215 (2001).
 - [23] P. Bak and K. Chen, Physica A **306**, 15 (2002).
 - [24] M. Paczusi and D. Hughes, Physica A, **342**, 158 (2004).
 - [25] P. Bak and M. Paczusi, Physica A, **348**, 277 (2005).
 - [26] E. G. Zweibel, Nature, **415**, 31 (2002).
 - [27] J. D. Gibbon and C. R. Doering, J. Fluid Mech., **478**, 227-235, 2003; *Intermittency and regularity issues in three-dimensional Navier-Stokes turbulence*, preprint (2003).
 - [28] H. Hentschel and I. Procaccia, Physica D **8**, 435 (1983).
 - [29] A. Kuo and S. Corrsin, J. Fluid Mech., **50**, 285, (1971).
 - [30] B. Mandelbrot, J. Fluid Mech. **62**, 331 (1974); *Turbulence and Navier-Stokes equations* (ed. R. Temam) Lect. Notes in Math. **565** (Springer, Berlin, 1976), pp 121.
 - [31] L. Caffarelli, R. Kohn and L. Nirenberg, Comm. Pure & Appl. Math., **35**, 771 (1982).
 - [32] P. Constantin and C. Foias, *Navier-Stokes Equations* (The University of Chicago Press, Chicago, 1988).
 - [33] C. Foias, O. Manley, R. Rosa and R. Temam, *Navier-Stokes equations and Turbulence* (Cambridge University Press, Cambridge, 2001).
 - [34] D. Sherrington and S. Kirkpatrick, Phys. Rev. Lett., **35** 1792 (1975).
 - [35] G. Parisi, M. Mézard and M. A. Virasoro, *Spin glass theory and beyond* (World Scientific, Singapore 1987).
 - [36] G. Parisi and F. Ricci-Tersenghi, J. Phys. A, **33**, 113 (2000).
 - [37] G. Hed, A. P. Young and E. Domany, Phys. Rev. Lett., **92**, 157201-1, (2004).
 - [38] D. A. Stariolo, Europhys. Lett., **55**, 726 (2001).
 - [39] S. Boettcher and M. Paczusi, Phys. Rev. E, **54**, 1082 (1996).
 - [40] The word ‘generic’ is being used to mean ‘typically’; of course the extreme case is that the integrand is positive, in which case no clusters form at all.
 - [41] As $p \rightarrow \infty$, $\langle \kappa_n^p \rangle^{1/p} \rightarrow \sup_{\Omega} \kappa_n$, which certainly lies within $\mathcal{A}_n^+$. The p -dependence of k_n^+ is suppressed.
 - [42] Since we expect $\mathcal{N}_n^+ \gg 1$, the estimate (14) implies that $L\lambda_{n+1}^+ \gg (\lambda_n^+)^2$. This is consistent with $\kappa_n > L^{-1}$ as in (1) but technically imposes an additional constraint.
 - [43] For instance, for a Fokker-Planck equation the Gibbs measure would be the most appropriate.